Proc. Indian Acad. Sci. (Math. Sci.) Vol. 113, No. 4, November 2003, pp. 379–393.
Printed in India

arXiv:math/0311523v1 [math.CA] 28 Nov 2003

Triebel–Lizorkin space estimates for multilinear operators of sublinear operators

LIU LANZHE

Department of Applied Mathematics, Hunan University, Changsha 410082,
People's Republic of China
Email: lanzheliu@263.net

MS received 23 March 2003

Abstract. In this paper, we obtain the continuity for some multilinear operators related to certain non-convolution operators on the Triebel–Lizorkin space. The operators include Littlewood–Paley operator and Marcinkiewicz operator.

Keywords. Multilinear operators; Triebel–Lizorkin space; Lipschitz space; Littlewood–Paley operator; Marcinkiewicz operator.

1. Introduction

Let T be the singular integral operator, a well-known result of Coifman, Rochberg and Weiss [6] which states that the commutator $[b,T]=T(bf)-bTf$ (where $b\in$ BMO) is bounded on $L^p(R^n)(1<p<\infty)$. Chanillo [1] proves a similar result when T is replaced by the fractional integral operator. In [9,11], these results on the Triebel–Lizorkin spaces and the case $b\in \mathrm{Lip}\beta$ (where $\mathrm{Lip}\beta$ is the homogeneous Lipschitz space) are obtained. The main purpose of this paper is to study the continuity for some multilinear operators related to certain convolution operators on the Triebel–Lizorkin spaces. In fact, we shall obtain the continuity on the Triebel–Lizorkin spaces for the multilinear operators only under certain conditions on the size of the operators. As applications, we prove the continuity of the multilinear operators related to the Littlewood–Paley operator and Marcinkiewicz operator on the Triebel–Lizorkin spaces.

2. Notations and results

In the sequel, Q will denote a cube of R^n with sides parallel to the axes, and for a cube Q, let $f_Q=|Q|^{-1}\int_Q f(x)\mathrm{d}x$ and $f^{\#}(x)=\sup_{x\in Q}|Q|^{-1}\int_Q|f(y)-f_Q|\mathrm{d}y$. For $1\le r<\infty$ and $0\le\delta$, let

$$M_{\delta,r}(f)(x)=\sup_{x\in Q}\left(\frac{1}{|Q|^{1-\delta r/n}}\int_Q|f(y)|^r\mathrm{d}y\right)^{1/r}.$$

We denote $M_{\delta,r}(f)=M_r(f)$ if $\delta=0$, which is the Hardy–Littlewood maximal function when $r=1$ (in this case, we denote $M_1(f)=M(f)$, see [12,13]). For $\beta>0$ and $p>1$, let

$\dot{F}_p^{\beta,\infty}$ be the homogeneous Triebel–Lizorkin space, the Lipschitz space $\dot{\wedge}_\beta$ is the space of functions f such that

$$\|f\|_{\dot{\wedge}_\beta} = \sup_{\substack{x,h\in R^n\\ h\neq 0}} |\Delta_h^{[\beta]+1} f(x)|/|h|^\beta < \infty,$$

where Δ_h^k denotes the kth difference operator [11].

We are going to consider the multilinear operator as follows:

Let m be a positive integer and A a function on R^n. We denote

$$R_{m+1}(A;x,y) = A(x) - \sum_{|\alpha|\le m} \frac{1}{\alpha!} D^\alpha A(y)(x-y)^\alpha.$$

DEFINITION 1.

Define $F(x,y,t)$ on $R^n \times R^n \times [0,+\infty)$. Then we denote

$$F_t(f)(x) = \int_{R^n} F(x,y,t) f(y)\mathrm{d}y$$

and

$$F_t^A(f)(x) = \int_{R^n} \frac{R_{m+1}(A;x,y)}{|x-y|^m} F(x,y,t) f(y)\mathrm{d}y.$$

Let H be the Hilbert space $H = \{h : \|h\| < \infty\}$. For each fixed $x \in R^n$, we view $F_t(f)(x)$ and $F_t^A(f)(x)$ as a mapping from $[0,+\infty)$ to H. Then, the multilinear operators related to F_t is defined by

$$T^A f(x) = \|F_t^A(f)(x)\|;$$

we also define that $Tf(x) = \|F_t f(x)\|$.

In particular, we consider the following two sublinear operators. Fix $\lambda > 1$.

DEFINITION 2.

Let $\varepsilon > 0$ and ψ be a fixed function which satisfies the following properties:

(1) $|\psi(x)| \le C(1+|x|)^{-(n+1)}$,
(2) $|\psi(x+y) - \psi(x)| \le C|y|^\varepsilon (1+|x|)^{-(n+1+\varepsilon)}$ when $2|y| < |x|$.

The multilinear Littlewood–Paley operator is defined by

$$g_\lambda^A(f)(x) = \left[\int\int_{R_+^{n+1}} \left(\frac{t}{t+|x-y|}\right)^{n\lambda} |F_t^A(f)(x,y)|^2 \frac{\mathrm{d}y\mathrm{d}t}{t^{n+1}}\right]^{1/2},$$

where

$$F_t^A(f)(x,y) = \int_{R^n} \frac{R_{m+1}(A;x,z)}{|x-z|^m} f(z)\psi_t(y-z)\mathrm{d}z$$

and $\psi_t(x) = t^{-n}\psi(x/t)$ for $t > 0$. We denote $F_t(f)(y) = f * \psi_t(y)$. We also define

$$g_\lambda(f)(x) = \left(\int\int_{R_+^{n+1}} \left(\frac{t}{t+|x-y|} \right)^{n\lambda} |F_t(f)(y)|^2 \frac{\mathrm{d}y\mathrm{d}t}{t^{n+1}} \right)^{1/2},$$

which is the Littlewood–Paley operator [13].

Let H be the Hilbert space $H = \{h : \|h\| = (\int\int_{R_+^{n+1}} |h(t)|^2 \mathrm{d}y\mathrm{d}t/t^{n+1})^{1/2} < \infty\}$. Then for each fixed $x \in R^n$, $F_t^A(f)(x,y)$ may be viewed as a mapping from $(0,+\infty)$ to H, and it is clear that

$$g_\lambda^A(f)(x) = \left\| \left(\frac{t}{t+|x-y|} \right)^{n\lambda/2} F_t^A(f)(x,y) \right\|,$$

$$g_\lambda(f)(x) = \left\| \left(\frac{t}{t+|x-y|} \right)^{n\lambda/2} F_t(f)(y) \right\|.$$

DEFINITION 3.

Let $0 < \gamma \le 1$ and Ω be a homogeneous of degree zero on R^n such that $\int_{S^{n-1}} \Omega(x')\mathrm{d}\sigma(x') = 0$. Assume that $\Omega \in \mathrm{Lip}_\gamma(S^{n-1})$, that is there exists a constant $M > 0$ such that for any $x,y \in S^{n-1}$, $|\Omega(x) - \Omega(y)| \le M|x-y|^\gamma$. We denote $\Gamma(x) = \{(y,t) \in R_+^{n+1} : |x-y| < t\}$ and the characteristic function of $\Gamma(x)$ by $\chi_{\Gamma(x)}$. The multilinear Marcinkiewicz integral operator is defined by

$$\mu_\lambda^A(f)(x) = \left[\int\int_{R_+^{n+1}} \left(\frac{t}{t+|x-y|} \right)^{n\lambda} |F_t^A(f)(x)|^2 \frac{\mathrm{d}y\mathrm{d}t}{t^{n+3}} \right]^{1/2},$$

where

$$F_t^A(f)(x,y) = \int_{|y-z|\le t} \frac{\Omega(y-z)}{|y-z|^{n-1}} \frac{R_{m+1}(A;x,z)}{|x-z|^m} f(z)\mathrm{d}z.$$

We denote that

$$F_t(f)(y) = \int_{|y-z|\le t} \frac{\Omega(y-z)}{|y-z|^{n-1}} f(z)\mathrm{d}z.$$

We also define that

$$\mu_\lambda(f)(x) = \left(\int\int_{R_+^{n+1}} \left(\frac{t}{t+|x-y|} \right)^{n\lambda} |F_t(f)(y)|^2 \frac{\mathrm{d}y\mathrm{d}t}{t^{n+3}} \right)^{1/2},$$

which is the Marcinkiewicz integral operator [14].

Let H be the Hilbert space $H = \{h : \|h\| = (\int\int_{R_+^{n+1}} |h(t)|^2 \mathrm{d}y\mathrm{d}t/t^{n+3})^{1/2} < \infty\}$. Then for each fixed $x \in R^n$, $F_t^A(f)(x,y)$ may be viewed as a mapping from $(0,+\infty)$ to H, and it is clear that

$$\mu_\lambda^A(f)(x) = \left\| \left(\frac{t}{t+|x-y|} \right)^{n\lambda/2} F_t^A(f)(x,y) \right\|,$$

$$\mu_\lambda(f)(x) = \left\| \left(\frac{t}{t+|x-y|} \right)^{n\lambda/2} F_t(f)(y) \right\|.$$

It is clear that Definitions 2 and 3 are the particular examples of Definition 1. Note that when $m = 0, T_A$ is just the commutator of F_t and A, and when $m > 0$, it is the non-trivial generalizations of the commutators. It is well-known that multilinear operators are of great interest in harmonic analysis and have been widely studied by many authors (see [2,3,4,5,7,8]). The main purpose of this paper is to study the continuity for the multilinear operators on the Triebel–Lizorkin spaces. We shall prove the following theorems in §3.

Theorem 1. *Let* g_λ^A *be the multilinear Littlewood–Paley operator as in Definition* 2*. If* $0 < \beta < 1/2$ *and* $D^\alpha A \in \dot{\wedge}_\beta$ *for* $|\alpha| = m$*. Then*

(a) g_λ^A *maps* $L^p(R^n)$ *continuously onto* $\dot{F}_p^{\beta,\infty}(R^n)$ *for* $1 < p < \infty$;
(b) g_λ^A *maps* $L^p(R^n)$ *continuously onto* $L^q(R^n)$ *for* $1 < p < n/\beta$ *and* $1/p - 1/q = \beta/n$.

Theorem 2. *Let* μ_λ^A *be the multilinear Marcinkiewicz operator as in Definition* 3*. If* $0 < \beta < 1/2$ *and* $D^\alpha A \in \dot{\wedge}_\beta$ *for* $|\alpha| = m$*. Then*

(a) μ_λ^A *maps* $L^p(R^n)$ *continuously onto* $\dot{F}_p^{\beta,\infty}(R^n)$ *for* $1 < p < \infty$;
(b) μ_λ^A *maps* $L^p(R^n)$ *continuously onto* $L^q(R^n)$ *for* $1 < p < n/\beta$ *and* $1/p - 1/q = \beta/n$.

3. Main theorem and proof

We first prove a general theorem.

Main Theorem. *Let* $0 < \beta < 1$ *and* $D^\alpha A \in \dot{\wedge}_\beta$ *for* $|\alpha| = m$*. Suppose* F_t, T *and* T^A *are the same as in Definition* 1*, if* T *is bounded on* $L^p(R^n)$ *for* $1 < r < \infty$ *and* T *satisfies the following size condition*:

$$\|F_t^A(f)(x) - F_t^A(f)(x_0)\| \le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\wedge}_\beta} |Q|^{\beta/n} M(f)(x)$$

for any cube Q *with* supp $f \subset (2Q)^c$ *and* $x \in Q$*. Then*

(a) T^A *maps* $L^p(R^n)$ *to* $\dot{F}_p^{\beta,\infty}(R^n)$ *for* $1 < p < \infty$;
(b) T^A *maps* $L^p(R^n)$ *to* $L^q(R^n)$ *for* $1 < p < n/\beta$ *and* $1/q = 1/p - \beta/n$.

To prove the theorem, we need the following lemmas.

Lemma 1.[11]. *For* $0 < \beta < 1, 1 < p < \infty$*, we have*

$$\|f\|_{\dot{F}_p^{\beta,\infty}} \approx \left\| \sup_Q \frac{1}{|Q|^{1+\beta/n}} \int_Q |f(x) - f_Q| \mathrm{d}x \right\|_{L^p}$$

$$\approx \left\| \sup_{\cdot \in Q} \inf_c \frac{1}{|Q|^{1+\beta/n}} \int_Q |f(x) - c| \mathrm{d}x \right\|_{L^p}.$$

Lemma 2.[11]. *For $0<\beta<1, 1\le p\le\infty$, we have*

$$\|f\|_{\dot\Lambda_\beta}\approx\sup_Q\frac{1}{|Q|^{1+\beta/n}}\int_Q|f(x)-f_Q|\mathrm{d}x$$

$$\approx\sup_Q\frac{1}{|Q|^{\beta/n}}\left(\frac{1}{|Q|}\int_Q|f(x)-f_Q|^p\mathrm{d}x\right)^{1/p}.$$

Lemma 3.[1,2]. *Suppose that $1\le r<p<n/\delta$ and $1/q=1/p-\delta/n$. Then*

$$\|M_{\delta,r}(f)\|_{L^q}\le C\|f\|_{L^p}.$$

Lemma 4.[5]. *Let A be a function on R^n and $D^\alpha A\in L^q(R^n)$ for $|\alpha|=m$ and some $q>n$. Then*

$$|R_m(A;x,y)|\le C|x-y|^m\sum_{|\alpha|=m}\left(\frac{1}{|\tilde Q(x,y)|}\int_{\tilde Q(x,y)}|D^\alpha A(z)|^q\mathrm{d}z\right)^{1/q},$$

where $\tilde Q(x,y)$ is the cube centered at x and having side length $5\sqrt{n}|x-y|$.

Proof of Main Theorem.
(a) Fix a cube $Q=Q(x_0,l)$ and $\tilde x\in Q$. Let $\tilde Q=5\sqrt{n}Q$ and $\tilde A(x)=A(x)-\sum_{|\alpha|=m}\frac{1}{\alpha!}(D^\alpha A)_{\tilde Q}x^\alpha$, then $R_m(A;x,y)=R_m(\tilde A;x,y)$ and $D^\alpha\tilde A=D^\alpha A-(D^\alpha A)_{\tilde Q}$ for $|\alpha|=m$. We write, for $f_1=f\chi_{\tilde Q}$ and $f_2=f\chi_{R^n\setminus\tilde Q}$,

$$\begin{aligned}F_t^A(f)(x)&=\int_{R^n}\frac{R_{m+1}(\tilde A;x,y)}{|x-y|^m}F(x,y,t)f(y)\mathrm{d}y\\&=\int_{R^n}\frac{R_{m+1}(\tilde A;x,y)}{|x-y|^m}F(x,y,t)f_2(y)\mathrm{d}y\\&\quad+\int_{R^n}\frac{R_m(\tilde A;x,y)}{|x-y|^m}F(x,y,t)f_1(y)\mathrm{d}y\\&\quad-\sum_{|\alpha|=m}\frac{1}{\alpha!}\int_{R^n}\frac{F(x,y,t)(x-y)^\alpha}{|x-y|^m}D^\alpha\tilde A(y)f_1(y)\mathrm{d}y,\end{aligned}$$

then

$$\begin{aligned}&|T^A(f)(x)-T^{\tilde A}(f_2)(x_0)|=|\|F_t^A(f)(x)\|-\|F_t^{\tilde A}(f_2)(x_0)\||\\&\le\left\|F_t\left(\frac{R_m(\tilde A;x,\cdot)}{|x-\cdot|^m}f_1\right)(x)\right\|+\sum_{|\alpha|=m}\frac{1}{\alpha!}\left\|F_t\left(\frac{(x-\cdot)^\alpha}{|x-\cdot|^m}D^\alpha\tilde A f_1\right)(x)\right\|\\&\quad+\|F_t^{\tilde A}(f_2)(x)-F_t^{\tilde A}(f_2)(x_0)\|=\mathrm{I}(x)+\mathrm{II}(x)+\mathrm{III}(x).\end{aligned}$$

Thus,

$$\begin{aligned}&\frac{1}{|Q|^{1+\beta/n}}\int_Q\left|T^A(f)(x)-T^{\tilde A}(f)(x_0)\right|\mathrm{d}x\\&\le\frac{1}{|Q|^{1+\beta/n}}\int_Q\mathrm{I}(x)\mathrm{d}x+\frac{1}{|Q|^{1+\beta/n}}\int_Q\mathrm{II}(x)\mathrm{d}x+\frac{1}{|Q|^{1+\beta/n}}\int_Q\mathrm{III}(x)\mathrm{d}x\\&:=\mathrm{I}+\mathrm{II}+\mathrm{III}.\end{aligned}$$

Now, let us estimate I, II and III, respectively. First, for $x \in Q$ and $y \in \tilde{Q}$, using Lemmas 2 and 4, we get

$$\begin{aligned} |R_m(\tilde{A};x,y)| &\le C|x-y|^m \sum_{|\alpha|=m} \sup_{x\in\tilde{Q}} |D^\alpha A(x) - (D^\alpha A)_{\tilde{Q}}| \\ &\le C|x-y|^m |Q|^{\beta/n} \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta}. \end{aligned}$$

Thus, by Holder's inequality and the L^r boundedness of T for $1 < r < p$, we obtain

$$\begin{aligned} \mathrm{I} &\le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} \frac{1}{|Q|} \int_Q |T(f_1)(x)| \mathrm{d}x \\ &\le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} \|T(f_1)\|_{L^r} |Q|^{-1/r} \\ &\le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} \|f_1\|_{L^r} |Q|^{-1/r} \\ &\le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} M_r(f)(\tilde{x}). \end{aligned}$$

Secondly, for $1 < r < q$, using the following inequality [11]:

$$\|D^\alpha A - (D^\alpha A)_{\tilde{Q}} f \chi_{\tilde{Q}}\|_{L^r} \le C|Q|^{1/r+\beta/n} \|D^\alpha A\|_{\dot{\Lambda}_\beta} M_r(f)(x)$$

and similar to the proof of I, we gain

$$\begin{aligned} \mathrm{II} &\le \frac{C}{|Q|^{1+\beta/n}} \sum_{|\alpha|=m} \|T((D^\alpha A - (D^\alpha A)_{\tilde{Q}}) f \chi_{\tilde{Q}})\|_{L^r} |Q|^{1-1/r} \\ &\le C|Q|^{-\beta/n-1/r} \sum_{|\alpha|=m} \|(D^\alpha A - (D^\alpha A)_{\tilde{Q}}) f \chi_{\tilde{Q}}\|_{L^r} \\ &\le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} M_r(f)(\tilde{x}). \end{aligned}$$

For III, using the size condition of T^A, we have

$$\mathrm{III} \le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} M(f)(\tilde{x}).$$

We now put these estimates together, and taking the supremum over all Q such that $\tilde{x} \in Q$, and using the L^p boundedness of M_r for $r < p$, we obtain

$$\|T^A(f)\|_{\dot{F}_p^{\beta,\infty}} \le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} \|f\|_{L^p}.$$

This completes the proof of (a).

(b) By same argument as in the proof of (a), we have

$$\begin{aligned} &\frac{1}{|Q|} \int_Q |T^A(f)(x) - T^{\tilde{A}}(f_2)(x_0)| \mathrm{d}x \\ &\quad \le C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\Lambda}_\beta} (M_{\beta,r}(f) + M_{\beta,1}(f)). \end{aligned}$$

Thus,

$$(T^A(f))^{\#} \leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\wedge}_\beta}(M_{\beta,r}(f)+M_{\beta,1}(f)).$$

Using Lemma 3, we gain

$$\begin{aligned}\|T^A(f)\|_{L^q} &\leq C\|(T^A(f))^{\#}\|_{L^q}\\ &\leq C \sum_{|\alpha|=m} \|D^\alpha A\|_{\dot{\wedge}_\beta}(\|M_{\beta,r}(f)\|_{L^q}+\|M_{\beta,1}(f)\|_{L^q}) \leq C\|f\|_{L^p}.\end{aligned}$$

This completes the proof of (b) and the Main Theorem.

To prove Theorems 1 and 2, we need the following lemma.

Lemma 5. *Let* $1<p<\infty$, $0<\beta<1$ *and* $D^\alpha A \in \dot{\wedge}_\beta$ *for* $|\alpha|=m$. *Then* g_λ^A *and* μ_λ^A *are all bounded on* $L^p(R^n)$.

Proof. For g_λ^A, by Minkowski inequality and the condition of ψ, we have

$$\begin{aligned}g_\lambda^A(f)(x) &\leq \int_{R^n} \frac{|f(z)||R_{m+1}(A;x,z)|}{|x-z|^m}\\ &\quad\times\left(\int_{R_+^{n+1}}\left(\frac{t}{t+|x-y|}\right)^{n\lambda}|\psi_t(y-z)|^2\frac{\mathrm{d}y\mathrm{d}t}{t^{1+n}}\right)^{1/2}\mathrm{d}z\\ &\leq C\int_{R^n} \frac{|f(z)||R_{m+1}(A;x,z)|}{|x-z|^m}\\ &\quad\times\left(\int_0^\infty\int_{R^n}\left(\frac{t}{t+|x-y|}\right)^{n\lambda}\frac{t^{-2n}}{(1+|y-z|/t)^{2n+2}}\right.\\ &\quad\times\left.\left(\frac{t}{t+|x-y|}\right)^{n\lambda}\frac{\mathrm{d}y\mathrm{d}t}{t^{1+n}}\right)^{1/2}\mathrm{d}z\\ &\leq C\int_{R^n} \frac{|f(z)||R_{m+1}(A;x,z)|}{|x-z|^m}\\ &\quad\times\left[\int_0^\infty\left(t^{-n}\int_{R^n}\left(\frac{t}{t+|x-y|}\right)^{n\lambda}\frac{\mathrm{d}y}{(t+|y-z|)^{2n+2}}\right)t\mathrm{d}t\right]^{1/2}\mathrm{d}z.\end{aligned}$$

Noting that

$$\begin{aligned}&t^{-n}\int_{R^n}\left(\frac{t}{t+|x-y|}\right)^{n\lambda}\frac{\mathrm{d}y}{(t+|y-z|)^{2n+2}}\\ &\quad\leq CM\left(\frac{1}{(t+|x-z|)^{2n+2}}\right) \leq \frac{C}{(t+|x-z|)^{2n+2}}\end{aligned}$$

and

$$\int_0^\infty \frac{t\mathrm{d}t}{(t+|x-z|)^{2n+2}} = C|x-z|^{-2n},$$

we obtain

$$\begin{aligned} g_\lambda^A(f)(x) &\le C\int_{R^n} \frac{|f(z)||R_{m+1}(A;x,z)|}{|x-z|^m}\left(\int_0^\infty \frac{t\mathrm{d}t}{(t+|x-z|)^{2n+2}}\right)^{1/2}\mathrm{dz} \\ &= C\int_{R^n}\frac{|f(z)||R_{m+1}(A;x,z)|}{|x-z|^{m+n}}\mathrm{dz}. \end{aligned}$$

For μ_λ, note that $|x-z|\le 2t$, $|y-z|\ge|x-z|-t\ge|x-z|-3t$ when $|x-y|\le t$, $|y-z|\le t$, and $|x-z|\le t(1+2^{k+1})\le 2^{k+2}t$, $|y-z|\ge|x-z|-2^{k+3}t$ when $|x-y|\le 2^{k+1}t$, $|y-z|\le t$, we have

$$\begin{aligned} \mu_\lambda^A(f)(x) \le{}& \int_{R^n}\Bigg[\int\int_{R_+^{n+1}}\left(\frac{t}{t+|x-y|}\right)^{n\lambda} \\ &\times\left(\frac{|\Omega(y-z)||R_{m+1}(A;x,z)||f(z)|}{|y-z|^{n-1}|x-z|^m}\right)^2\chi_{\Gamma(z)}(y,t)\frac{\mathrm{d}y\mathrm{d}t}{t^{n+3}}\Bigg]^{1/2}\mathrm{dz} \\ \le{}& C\int_{R^n}\frac{|R_{m+1}(A;x,z)||f(z)|}{|x-z|^m} \\ &\times\left[\int_0^\infty\int_{|x-y|\le t}\left(\frac{t}{t+|x-y|}\right)^{n\lambda}\frac{\chi_{\Gamma(z)}(y,t)}{(|x-z|-3t)^{2n-2}}\frac{\mathrm{d}y\mathrm{d}t}{t^{n+3}}\right]^{1/2}\mathrm{dz} \\ &+C\int_{R^n}\frac{|R_{m+1}(A;x,z)||f(z)|}{|x-z|^m} \\ &\times\left[\int_0^\infty\sum_{k=0}^\infty\int_{2^kt<|x-y|\le 2^{k+1}t}\left(\frac{t}{t+|x-y|}\right)^{n\lambda}\frac{\chi_{\Gamma(z)}(y,t)t^{-n-3}\mathrm{d}y\mathrm{d}t}{(|x-z|-2^{k+3}t)^{2n-2}}\right]^{1/2}\mathrm{dz} \\ \le{}& C\int_{R_n}\frac{|R_{m+1}(A;x,z)||f(z)|}{|x-z|^{m+1/2}}\left[\int_{|x-z|/2}^\infty\frac{\mathrm{d}t}{(|x-z|-3t)^{2n}}\right]^{1/2}\mathrm{dz} \\ &+C\int_{R_n}\frac{|R_{m+1}(A;x,z)||f(z)|}{|x-z|^{m+1/2}} \\ &\times\left[\sum_{k=0}^\infty\int_{2^{-2-k}|x-z|}^\infty 2^{-kn\lambda}(2^kt)^nt^{-n}\frac{2^k\mathrm{d}t}{(|x-z|-2^{k+3}t)^{2n}}\right]^{1/2}\mathrm{dz} \\ \le{}& C\int_{R_n}\frac{|R_{m+1}(A;x,z)||f(z)|}{|x-z|^{m+n}}\mathrm{dz} \\ &+C\int_{R_n}\frac{|R_{m+1}(A;x,z)||f(z)|}{|x-z|^{m+n}}\mathrm{dz}\left[\sum_{k=0}^\infty 2^{kn(1-\lambda)}\right]^{1/2} \\ ={}& C\int_{R_n}\frac{|R_{m+1}(A;x,z)|}{|x-z|^{m+n}}|f(z)|\mathrm{dz}. \end{aligned}$$

Thus, the lemma follows from [2].

Now we can prove Theorems 1 and 2. Since g_λ and μ_λ are all bounded on $L^p(R^n)$ for $1 < p < \infty$ by Lemma 5, it suffices to verify that g_λ^A and μ_λ^A satisfy the size condition in the Main Theorem.

For g_λ^A, we write

$$\begin{aligned}
F_t^A(f)(x,y) &= \int \frac{R_{m+1}(\tilde{A};x,z)}{|x-z|^m}\psi_t(y-z)f(z)\mathrm{d}z \\
&= \int \frac{R_{m+1}(\tilde{A};x,z)}{|x-z|^m}\psi_t(y-z)f_2(z)\mathrm{d}z \\
&\quad + \int \frac{R_m(\tilde{A};x,z)}{|x-z|^m}\psi_t(y-z)f_1(z)\mathrm{d}z \\
&\quad - \sum_{|\alpha|=m}\frac{1}{\alpha!}\int \frac{(x-z)^\alpha \psi_t(y-z)}{|x-z|^m}D^\alpha \tilde{A}(z)f_1(z)\mathrm{d}z,
\end{aligned}$$

then

$$\begin{aligned}
&\frac{1}{|Q|^{1+\beta/n}}\int_Q \left|g_\lambda^A(f)(x) - g_\lambda^{\tilde{A}}(f_2)(x_0)\right|\mathrm{d}x \\
&= \frac{1}{|Q|^{1+\beta/n}}\int_Q \Bigg|\left\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^A(f)(x,y)\right\| \\
&\quad - \left\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f)(x_0,y)\right\|\Bigg|\mathrm{d}x \\
&\le \frac{1}{|Q|^{1+\beta/n}}\int_Q \left\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t\left(\frac{R_m(\tilde{A};x,\cdot)}{|x-\cdot|^m}f_1\right)(y)\right\|\mathrm{d}x \\
&\quad + \frac{1}{|Q|^{1+\beta/n}}\int_Q \sum_{|\alpha|=m}\frac{1}{\alpha!}\Bigg\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2} \\
&\quad \times F_t\left(\frac{(x-\cdot)^\alpha}{|x-\cdot|^m}D^\alpha \tilde{A} f_1\right)(y)\Bigg\|\mathrm{d}x \\
&\quad + \frac{1}{|Q|^{1+\beta/n}}\int_Q \Bigg\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x,y) \\
&\quad - \left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x_0,y)\Bigg\|\mathrm{d}x \\
&:= \mathrm{I} + \mathrm{II} + \mathrm{III}.
\end{aligned}$$

For I and II, similar to the proof of Lemma 5 and the Main Theorem, we get

$$\begin{aligned}
\mathrm{I} &\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\frac{1}{|Q|}\int_Q |g_\lambda(f_1)(x)|\mathrm{d}x \\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\|g_\lambda(f_1)\|_{L^r}|Q|^{-1/r}
\end{aligned}$$

$$
\begin{aligned}
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot\Lambda_\beta}\|f_1\|_{L^r}|Q|^{-1/r}\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot\Lambda_\beta}M_r(f)(\tilde{x})
\end{aligned}
$$

and

$$
\begin{aligned}
\mathrm{II} &\le \frac{C}{|Q|^{1+\beta/n}}\sum_{|\alpha|=m}\|g_\lambda((D^\alpha A-(D^\alpha A)_{\tilde{Q}})f\chi_{\tilde{Q}})\|_{L^r}|Q|^{1-1/r}\\
&\le C|Q|^{-\beta/n-1/r}\sum_{|\alpha|=m}\|(D^\alpha A-(D^\alpha A)_{\tilde{Q}})f\chi_{\tilde{Q}}\|_{L^r}\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot\Lambda_\beta}M_r(f)(\tilde{x}).
\end{aligned}
$$

For III, we write

$$
\begin{aligned}
&\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x,y)-\left(\frac{t}{t+|x_0-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x_0,y)\\
&=\int_{R^n}\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}\left[\frac{1}{|x-z|^m}-\frac{1}{|x_0-z|^m}\right]\\
&\quad\times R_m(\tilde{A};x,z)\psi_t(y-z)f_2(z)\mathrm{d}z\\
&\quad+\int_{R^n}\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}\frac{\psi_t(y-z)f_2(z)}{|x_0-z|^m}[R_m(\tilde{A};x,z)-R_m(\tilde{A};x_0,z)]\mathrm{d}z\\
&\quad+\int_{R^n}\left[\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}-\left(\frac{t}{t+|x_0-y|}\right)^{n\lambda/2}\right]\\
&\quad\times\frac{R_m(\tilde{A};x_0,z)\psi_t(y-z)f_2(z)}{|x_0-z|^m}\mathrm{d}z\\
&\quad-\sum_{|\alpha|=m}\frac{1}{\alpha!}\int_{R^n}\left[\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}\frac{(x-z)^\alpha}{|x-z|^m}\right.\\
&\quad\left.-\left(\frac{t}{t+|x_0-y|}\right)^{n\lambda/2}\frac{(x_0-z)^\alpha}{|x_0-z|^m}\right]D^\alpha\tilde{A}(z)\psi_t(y-z)f_2(z)\mathrm{d}z\\
&:=\mathrm{III}_1+\mathrm{III}_2+\mathrm{III}_3+\mathrm{III}_4.
\end{aligned}
$$

Note that $|x-z|\sim|x_0-z|$ for $x\in Q$ and $z\in R^n\setminus\tilde{Q}$. By the condition of ψ and similar to the proof of Lemma 5, we obtain

$$
\begin{aligned}
&\frac{1}{|Q|^{1+\beta/n}}\int_Q\|\mathrm{III}_1\|\mathrm{d}x\\
&\le\frac{C}{|Q|^{1+\beta/n}}\int_Q\left(\int_{R^n\setminus\tilde{Q}}\frac{|x-x_0|}{|x_0-z|^{m+n+1}}|R_m(\tilde{A};x,z)||f(z)|\mathrm{d}z\right)\mathrm{d}x
\end{aligned}
$$

$$\le C\sum_{|\alpha|=m}\|D^{\alpha}A\|_{\dot{\Lambda}_{\beta}}\sum_{k=0}^{\infty}\int_{2^{k+1}\tilde{Q}\setminus 2^{k+1}\tilde{Q}}\frac{|x-x_0|}{|x_0-z|^{n+1}}|f(z)|\mathrm{d}z$$

$$\le C\sum_{|\alpha|=m}\|D^{\alpha}A\|_{\dot{\Lambda}_{\beta}}\sum_{k=1}^{\infty}2^{-k}\frac{1}{|2^k\tilde{Q}|}\int_{2^k\tilde{Q}}|f(z)|\mathrm{d}z$$

$$\le C\sum_{|\alpha|=m}\|D^{\alpha}A\|_{\dot{\Lambda}_{\beta}}\sum_{k=1}^{\infty}2^{-k}M(f)(\tilde{x})$$

$$\le C\sum_{|\alpha|=m}\|D^{\alpha}A\|_{\dot{\Lambda}_{\beta}}M(f)(\tilde{x}).$$

For III$_2$, by the formula [5]

$$R_m(\tilde{A};x,z)-R_m(\tilde{A};x_0,z)=\sum_{|\eta|<m}\frac{1}{\eta!}R_{m-|\eta|}(D^{\eta}\tilde{A};x,x_0)(x-z)^{\eta}$$

and Lemma 4, we get

$$|R_m(\tilde{A};x,z)-R_m(\tilde{A};x_0,z)|$$
$$\le C\sum_{|\alpha|=m}\|D^{\alpha}A\|_{\dot{\Lambda}_{\beta}}|Q|^{\beta/n}|x-x_0||x_0-z|^{m-1}.$$

Thus

$$\frac{1}{|Q|^{1+\beta/n}}\int_Q\|\mathrm{III}_2\|\mathrm{d}x$$

$$\le C\frac{1}{|Q|^{1+\beta/n}}\int_Q\int_{R^n\setminus\tilde{Q}}\frac{|R_m(\tilde{A};x,z)-R_m(\tilde{A};x_0,z)|}{|x_0-z|^{m+n}}|f(z)|\mathrm{d}z\mathrm{d}x$$

$$\le C\sum_{|\alpha|=m}\|D^{\alpha}A\|_{\dot{\Lambda}_{\beta}}\sum_{k=0}^{\infty}\int_{2^{k+1}\tilde{Q}\setminus 2^k\tilde{Q}}\frac{|x-x_0|}{|x_0-y|^{n+1}}|f(z)|\mathrm{d}z$$

$$\le C\sum_{|\alpha|=m}\|D^{\alpha}A\|_{\dot{\Lambda}_{\beta}}|Q|^{\beta/n}M(f)(\tilde{x}).$$

For III$_3$, by the inequality: $a^{1/2}-b^{1/2}\le(a-b)^{1/2}$ for $a\ge b>0$, we gain

$$\frac{1}{|Q|^{1+\beta/n}}\int_Q\|\mathrm{III}_3\|\mathrm{d}x\le\frac{C}{|Q|^{1+\beta/n}}$$

$$\times\int_Q\int_{R^n}\left(\int_{R^{n+1}_{+}}\left[\frac{t^{n\lambda/2}|x-x_0|^{1/2}|\psi_t(y-z)||R_m(\tilde{A};x_0,z)||f_2(z)|}{|x-z|^m(t+|x-y|)^{(n\lambda+1)/2}}\right]^2\frac{\mathrm{d}y\mathrm{d}t}{t^{n+1}}\right)^{1/2}\mathrm{d}z\mathrm{d}x$$

$$
\begin{aligned}
&\le \frac{C}{|Q|^{1+\beta/n}}\int_Q\int_{R^n}\frac{|f_2(z)||R_m(\tilde{A};x_0,z)||x-x_0|^{1/2}}{|x-z|^m}\\
&\qquad\times\left(\int_0^\infty\frac{\mathrm{d}t}{(t+|x-z|)^{2n+2}}\right)^{1/2}\mathrm{d}z\mathrm{d}x\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\int_{R^n}\frac{|x-x_0|^{1/2}}{|x_0-z|^{n+1/2}}|f_2(z)|\mathrm{d}z\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M(f)(\tilde{x}).
\end{aligned}
$$

For III_4, by Lemma 4, we know that $|D^\alpha A(z)-(D^\alpha A)_{\tilde{Q}}|\le\|D^\alpha A\|_{\dot{\Lambda}_\beta}|x_0-z|^\beta$. Thus, similar to the proof of III_1 and III_3, we obtain

$$
\begin{aligned}
&\frac{1}{|Q|^{1+\beta/n}}\int_Q\|\mathrm{III}_4\|\mathrm{d}x\\
&\le\frac{C}{|Q|^{1+\beta/n}}\int_Q\sum_{|\alpha|=m}\int_{R^n}\left(\frac{|x-x_0|}{|x_0-z|^{n+1}}+\frac{|x-x_0|^{1/2}}{|x_0-z|^{n+1/2}}\right)\\
&\qquad\times|f_2(z)||D^\alpha\tilde{A}(z)|\mathrm{d}z\mathrm{d}x\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\sum_{k=1}^\infty(2^{k(\beta-1)}+2^{k(\beta-1/2)})M(f)(\tilde{x})\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M(f)(\tilde{x}).
\end{aligned}
$$

Thus,

$$\mathrm{III}\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M(f)(\tilde{x}).$$

For μ_λ^A, similarly, we have

$$
\begin{aligned}
&\frac{1}{|Q|^{1+\beta/n}}\int_Q|\mu_\lambda^A(f)(x)-\mu_\lambda^{\tilde{A}}(f_2)(x_0)|\mathrm{d}x\\
&=\frac{1}{|Q|^{1+\beta/n}}\int_Q\left|\left\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^A(f)(x,y)\right\|\right.\\
&\qquad\left.-\left\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x_0,y)\right\|\right|\mathrm{d}x\\
&\le\frac{1}{|Q|^{1+\beta/n}}\int_Q\left\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t\left(\frac{R_m(\tilde{A};x,\cdot)}{|x-\cdot|^m}f_1\right)(y)\right\|\mathrm{d}x\\
&\qquad+\frac{1}{|Q|^{1+\beta/n}}\int_Q\sum_{|\alpha|=m}\frac{1}{\alpha!}\left\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t\left(\frac{(x-\cdot)^\alpha}{|x-\cdot|^m}D^\alpha\tilde{A}f_1\right)(y)\right\|\mathrm{d}x
\end{aligned}
$$

$$
\begin{aligned}
&+\frac{1}{|Q|^{1+\beta/n}}\int_Q\left\|\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x,y)\right.\\
&\left.-\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x_0,y)\right\|\mathrm{d}x\\
&:=\mathrm{I}+\mathrm{II}+\mathrm{III}
\end{aligned}
$$

and

$$
\begin{aligned}
\mathrm{I}&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\frac{1}{|Q|}\int_Q|\mu_\lambda(f_1)(x)|\mathrm{d}x\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\|\mu_\lambda(f_1)\|_{L^r}|Q|^{-1/r}\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\|f_1\|_{L^r}|Q|^{-1/r}\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M_r(f)(\tilde{x}),\\
\mathrm{II}&\le\frac{C}{|Q|^{1+\beta/n}}\sum_{|\alpha|=m}\|\mu_\lambda((D^\alpha A-(D^\alpha A)_{\tilde{Q}})f\chi_{\tilde{Q}})\|_{L^r}|Q|^{1-1/r}\\
&\le C|Q|^{-\beta/n-1/r}\sum_{|\alpha|=m}\|(D^\alpha A-(D^\alpha A)_{\tilde{Q}})f\chi_{\tilde{Q}}\|_{L^r}\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M_r(f)(\tilde{x}).
\end{aligned}
$$

For III, we write

$$
\begin{aligned}
&\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x,y)-\left(\frac{t}{t+|x_0-y|}\right)^{n\lambda/2}F_t^{\tilde{A}}(f_2)(x_0,y)\\
&=\int_{|y-z|\le t}\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}\left[\frac{1}{|x-z|^m}-\frac{1}{|x_0-z|^m}\right]\\
&\quad\times\frac{\Omega(y-z)R_m(\tilde{A};x,z)f_2(z)}{|y-z|^{n-1}}\mathrm{d}z\\
&\quad+\int_{|y-z|\le t}\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}\frac{\Omega(y-z)f_2(z)}{|y-z|^{n-1}|x_0-z|^m}\\
&\quad\times[R_m(\tilde{A};x,z)-R_m(\tilde{A};x_0,z)]\mathrm{d}z\\
&\quad+\int_{|y-z|\le t}\left[\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}-\left(\frac{t}{t+|x_0-y|}\right)^{n\lambda/2}\right]\\
&\quad\times\frac{\Omega(y-z)R_m(\tilde{A};x_0,z)f_2(z)}{|y-z|^{n-1}|x_0-z|^m}\mathrm{d}z
\end{aligned}
$$

$$
\begin{aligned}
&-\sum_{|\alpha|=m}\frac{1}{\alpha!}\int_{|y-z|\le t}\left[\left(\frac{t}{t+|x-y|}\right)^{n\lambda/2}\frac{(x-z)^\alpha}{|x-z|^m}\right.\\
&\left.-\left(\frac{t}{t+|x_0-y|}\right)^{n\lambda/2}\frac{(x_0-z)^\alpha}{|x_0-z|^m}\right]\frac{\Omega(y-z)D^\alpha\tilde{A}(z)f_2(z)}{|y-z|^{n-1}}\mathrm{d}z\\
&:=\mathrm{III}_1+\mathrm{III}_2+\mathrm{III}_3+\mathrm{III}_4.
\end{aligned}
$$

By the condition of Ω and similar to the proof of Lemma 5, we get

$$
\begin{aligned}
&\frac{1}{|Q|^{1+\beta/n}}\int_Q\|\mathrm{III}_1\|\mathrm{d}x\\
&\le\frac{C}{|Q|^{1+\beta/n}}\int_Q\left(\int_{R^n\backslash\tilde{Q}}\frac{|x-x_0|}{|x_0-z|^{m+n+1}}|R_m(\tilde{A};x,z)\|f(z)|\mathrm{d}z\right)\mathrm{d}x\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\sum_{k=0}^{\infty}\int_{2^{k+1}\tilde{Q}\backslash 2^{k+1}\tilde{Q}}\frac{|x-x_0|}{|x_0-z|^{n+1}}|f(z)|\mathrm{d}z\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M(f)(\tilde{x}),\\
&\frac{1}{|Q|^{1+\beta/n}}\int_Q\|\mathrm{III}_2\|\mathrm{d}x\\
&\le C\frac{1}{|Q|^{1+\beta/n}}\int_Q\int_{R^n\backslash\tilde{Q}}\frac{|R_m(\tilde{A};x,z)-R_m(\tilde{A};x_0,z)|}{|x_0-z|^{m+n}}|f(z)|\mathrm{d}z\mathrm{d}x\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\sum_{k=0}^{\infty}\int_{2^{k+1}\tilde{Q}\backslash 2^{k}\tilde{Q}}\frac{|x-x_0|}{|x_0-y|^{n+1}}|f(z)|\mathrm{d}z\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}|Q|^{\beta/n}M(f)(\tilde{x}),\\
&\frac{1}{|Q|^{1+\beta/n}}\int_Q\|\mathrm{III}_3\|\mathrm{d}x\\
&\le\frac{C}{|Q|^{1+\beta/n}}\int_Q\int_{R^n}\left(\int_{R^{n+1}_+}\left[\frac{t^{n\lambda/2}|x-x_0|^{1/2}\chi_{\Gamma(z)}(y,t)|f_2(z)|}{(t+|x-y|)^{(n\lambda+1)/2}|y-z|^{n-1}}\right.\right.\\
&\qquad\left.\left.\times\frac{|R_m(\tilde{A};x_0,z)|}{|x_0-z|^m}\right]^2\frac{\mathrm{d}y\mathrm{d}t}{t^{n+3}}\right)^{1/2}\mathrm{d}z\mathrm{d}x\\
&\le\frac{C}{|Q|^{1+\beta/n}}\int_Q\int_{R^n}\frac{|R_m(\tilde{A};x_0,z)||f_2(z)||x-x_0|^{1/2}}{|x_0-z|^{m+n+1/2}}\mathrm{d}z\mathrm{d}x\\
&\le C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M(f)(\tilde{x})
\end{aligned}
$$

and

$$\begin{aligned}
&\frac{1}{|Q|^{1+\beta/n}}\int_Q \|\mathrm{III}_4\|\mathrm{d}x \\
&\leq \frac{C}{|Q|^{1+\beta/n}}\int_Q \sum_{|\alpha|=m}\int_{R^n}\left(\frac{|x-x_0|}{|x_0-z|^{n+1}}+\frac{|x-x_0|^{1/2}}{|x_0-z|^{n+1/2}}\right) \\
&\quad \times |f_2(z)||D^\alpha \tilde{A}(z)|\mathrm{d}z\mathrm{d}x \\
&\leq C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}\sum_{k=1}^{\infty}(2^{k(\beta-1)}+2^{k(\beta-1/2)})M(f)(\tilde{x}) \\
&\leq C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M(f)(\tilde{x}).
\end{aligned}$$

Thus,

$$\mathrm{III} \leq C\sum_{|\alpha|=m}\|D^\alpha A\|_{\dot{\Lambda}_\beta}M(f)(\tilde{x}).$$

These yield the desired results.

References

[1] Chanillo S, A note on commutators, *Indiana Univ. Math. J.* **31** (1982) 7–16

[2] Chen W G, Besov estimates for a class of multilinear singular integrals, *Acta Math. Sinica* **16** (2000) 613–626

[3] Cohen J, A sharp estimate for a multilinear singular integral on R^n, *Indiana Univ. Math. J.* **30** (1981) 693–702

[4] Cohen J and Gosselin J, On multilinear singular integral operators on R^n, *Studia Math.* **72** (1982) 199–223

[5] Cohen J and Gosselin J, A BMO estimate for multilinear singular integral operators, *Illinois J. Math.* **30** (1986) 445–465

[6] Coifman R, Rochberg R and Weiss G, Factorization theorems for Hardy spaces in several variables, *Ann. Math.* **103** (1976) 611–635

[7] Ding Y, A note on multilinear fractional integrals with rough kernel, *Adv. Math. (China)* **30** (2001) 238–246.

[8] Ding Y and Lu S Z, Weighted boundedness for a class rough multilinear operators, *Acta Math. Sinica* **3** (2001) 517–526

[9] Janson S, Mean oscillation and commutators of singular integral operators, *Ark. Math.* **16** (1978) 263–270

[10] Liu Lanzhe, Continuity for commutators of Littlewood–Paley operator on certain Hardy spaces, *J. Korean Math. Soc.* **40(1)** (2003) 41–60

[11] Paluszynski M, Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss, *Indiana Univ. Math. J.* **44** (1995) 1–17

[12] Stein E M, Harmonic analysis: Real variable methods, orthogonality and oscillatory integrals (Princeton, NJ: Princeton Univ. Press) (1993)

[13] Torchnisky A, The real variable methods in harmonic analysis, *Pure Appl. Math.* (New York: Academic Press) (1986) vol. 123

[14] Torchnisky A and Wang S, A note on the Marcinkiewicz integral, *Colloq. Math.* **60/61** (1990) 235–243